

The properties of Galactic globular cluster sub-systems

A. D. Mackey¹ and Sidney van den Bergh²

¹*Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK. E-mail: dmackey@ast.cam.ac.uk*

²*Dominion Astrophysical Observatory, Herzberg Institute of Astrophysics, National Research Council of Canada, 5071 West Saanich Road, Victoria, British Columbia, V9E 2E7, Canada. E-mail: sidney.vandenbergh@nrc-cnrc.gc.ca*

Accepted – Received –

ABSTRACT

In this paper we compare the properties of three sub-systems of Galactic globular clusters, which are defined according to metallicity and horizontal branch morphology. We specifically focus on cluster luminosities, structures, surface brightnesses and ellipticities. It is shown that the so-called “young” halo (YH) clusters, which are thought to have formed in external satellite galaxies, exhibit characteristics which are clearly distinct from those of the “old” halo (OH) and bulge/disk (BD) clusters, the majority of which are believed to be Galactic natives. The properties of the YH objects are, in many respects, similar to those of clusters belonging to a number of present day satellite dwarf galaxies. The OH and BD populations have apparently been strongly modified by destructive tidal forces and shocks in the inner Galaxy. By comparing the properties of the three cluster sub-systems, we estimate that the present population of native Galactic clusters may only represent approximately two-thirds of the original population. Several clusters with low surface brightnesses are observed to be highly flattened. We briefly speculate on the possibility that this ellipticity reflects the intrinsic flattening of dark matter mini-halos in which these optically dim clusters might be embedded. Finally, we examine the distribution of clusters on the size ($\log R_h$) vs. luminosity (M_V) plane. Three objects are seen to fall well above the sharp upper envelope of the main distribution of clusters on the size-luminosity plane: ω Centauri, M54, and NGC 2419. All three of these objects have previously, and independently, been suggested to be the stripped cores of former dwarf galaxies. This suspicion is strengthened by the additional observation that the massive cluster G1 in M31, plus a number of the most luminous clusters in NGC 5128 also fall in the same region of the $\log R_h$ vs. M_V plane. All of the latter objects have previously been suggested as the stripped cores of now defunct dwarf galaxies.

Key words: Galaxy: halo, formation – globular clusters: general

1 INTRODUCTION

Our Galaxy contains ~ 150 globular clusters, located at distances between 500 pc and 120 kpc from the Galactic centre. Because they are generally bright, and probe such a large range of Galactocentric radii, these objects constitute important tools for both tracing the properties of our Galaxy in the present epoch and for piecing together its formation history. They are also interesting objects in their own right, since they have the potential to provide information on star and cluster formation processes. Furthermore they provide useful laboratories for the study of self-gravitating stellar systems.

As a globular cluster evolves dynamically its core contracts and its envelope expands. However, N -body calculations (e.g., Spitzer & Thuan (1972); Lightman & Shapiro (1978); Aarseth & Heggie (1998)) show that R_h – the radius that contains half of the cluster stars in projection, generally changes very little over periods as long as ten relaxation times. This suggests that R_h , like $[\text{Fe}/\text{H}]$, may provide valuable information on physical conditions during the era of globular cluster formation. However, an important caveat

is that metallicity dependent differences of up to 20 per cent can occur between the half-light radii and the half-mass radii of clusters of differing metallicity (Jordán 2004). The reason for this metallicity dependence is that the masses of the most luminous surviving stars become larger as the cluster metallicity increases. As a result the density profile of a metal-rich cluster will be slightly more centrally concentrated than that of an otherwise similar metal-poor cluster. Nevertheless, the half-light radii of clusters are of considerable interest because they are stable over many relaxation times.

The work of Mackey & Gilmore (2004) has recently re-emphasized the fact that Galactic globular clusters comprise a very inhomogeneous class of objects. It is therefore useful to try to identify sub-groups made up of clusters having similar characteristics. One of the most successful classification schemes for the Galactic globular clusters is that introduced over a decade ago (see e.g., Zinn (1993); van den Bergh (1993), and references therein), in which clusters are grouped according to both their metallicity and their horizontal-branch (HB) morphology. These parameters are closely related to a number of other cluster properties such as, for example,

Galactocentric radius and age. Such a division of globular clusters can therefore provide us with a great deal of information about the evolutionary history of the Galaxy. Recently, Mackey & Gilmore (2004) updated the original classification scheme to cover the full Galactic globular cluster sample for which information is presently available. By comparing the different Galactic sub-groups to the cluster systems in nearby dwarf galaxies (i.e., the LMC, SMC, and Fornax and Sagittarius dwarf spheroidal galaxies) these authors confirmed that ~ 30 per cent of the Galactic globular cluster system possesses properties consistent with an origin external to the Milky Way. In other words, such clusters are most likely to have formed within much smaller galaxies (perhaps similar to the Local Group dwarf spheroidals observed at the present epoch) which later merged with the Milky Way system. Furthermore, Mackey & Gilmore (2004) showed that classifying Galactic globular clusters by metallicity and HB morphology leads quite naturally to a segregation of clusters by size. These authors used cluster core radius R_c as their size diagnostic; however, unlike the half-light radius R_h , this parameter is quite sensitive to the dynamically evolving state of a globular cluster. It is therefore of interest to extend the analysis of Mackey & Gilmore (2004) to additional parameters, such as R_h , which “remember” more about the initial sizes of clusters, and hence about physical conditions in the Galaxy and/or in its progenitors, at early epochs.

In the present paper we shall examine the different Galactic globular cluster sub-systems in terms of cluster structures, luminosities, and surface brightnesses. Section 2 lists the relevant data for each of the Galactic globular clusters and presents the classification of each object according to Mackey & Gilmore (2004). In Section 3 we use these data to investigate the collective properties of each cluster sub-system, while in Section 4 we discuss the possible links between half-light radius and luminosity both for Galactic globular clusters and for clusters belonging to some other nearby Local Group dwarf galaxies. Finally, we discuss what these parameters might be able to tell us about the origins and evolution of different types of clusters.

2 CLUSTER DATA AND CLASSIFICATIONS

In Table 1 we list all 150 known Galactic globular clusters, including the handful of clusters that are thought to be associated with the Sagittarius dwarf galaxy. The structures of most of these objects have been measured. For each such cluster we list the half-light radius (R_h) and tidal radius (R_t). We have adopted the majority of these values from the online database of Harris (1996) (2003 update). Since this compilation lists R_h and R_t as angular diameters, we have converted them to parsecs using the cluster distances that are also listed by Harris. For a few clusters the database does not have entries for R_h and R_t . We have been able to fill in these missing values for four clusters: IC 1257 (Côté 1999), BH 176, Terzan 4, and 1636-283 (Webbink 1985). In addition we have, for HP 1, adopted the measurements of Ortolani, Bica & Barbuy (1997), who used a full colour-magnitude diagram to derive a distance less than half that listed by Harris, in good agreement with the measurement by Davidge (2000). Also given in Table 1 are the Galactocentric distance R_{gc} , the integrated luminosity M_V , and (where available) the ellipticity ($\epsilon = 1 - b/a$, where a is the cluster semi-major axis and b the semi-minor axis), again all taken from the Harris database.

Using the half-light radius and integrated luminosity for each cluster, we have calculated I_h , the half-light intensity. This is de-

fined as $I_h = \frac{L}{2\pi R_h^2}$, and is measured in units of solar luminosities per square parsec ($L_\odot \text{ pc}^{-2}$). It is essentially a measure of the mean surface brightness of a cluster within R_h , but in terms of absolute units rather than observational units. Because of the evaporation of stars from dynamically evolving clusters, I_h is more sensitive to evolutionary effects than is R_h . Nevertheless, this parameter turns out to be useful when examining cluster classifications. Finally, Table 1 lists the metallicity $[\text{Fe}/\text{H}]$ and HB morphology index of each cluster, along with the formal type classification from Mackey & Gilmore (2004). As noted in Section 1, this classification scheme follows closely those of Zinn (1993) and of van den Bergh (1993) (see also the references therein). To briefly review this scheme, the clusters are grouped by metallicity and HB-type*. The metal rich globular clusters with $[\text{Fe}/\text{H}] > -0.8$ all have red horizontal branches and are spatially confined to the bulge and inner disk of the Galaxy. Clusters in this sub-group are designated bulge/disk (BD) objects. In contrast, the metal-poor clusters (with $[\text{Fe}/\text{H}] \leq -0.8$) are seen to have a wide range in HB-type and are generally situated in the Galactic halo.

Most of the inner halo clusters exhibit quite a tight relationship between HB-type and $[\text{Fe}/\text{H}]$. On the other hand many of the outer halo clusters have much redder HB-types at given metallicity (this is the classical “second-parameter phenomenon”, where a parameter in addition to metallicity affects HB morphology). Following Zinn (1993), the “old” halo (OH) and “young” halo (YH) sub-systems are defined by measuring the offset in HB-type at given $[\text{Fe}/\text{H}]$ of each cluster from the fiducial line which traces the relationship between HB-type and $[\text{Fe}/\text{H}]$ for the inner halo clusters†. Objects for which the offset in HB-type is greater than -0.3 (i.e., objects which do not lie far to the red of the fiducial line) constitute the OH class, while those with offset less than -0.3 (i.e., clusters which have red HB-type at given $[\text{Fe}/\text{H}]$) form the YH class.

The majority of the metallicity and HB-type parameters we list in Table 1 are taken from the Harris database, except where they were updated by Mackey & Gilmore (2004) (one metallicity value and 28 HB indices). There are five classification types listed in Table 1: BD for the bulge/disk clusters (37 objects); OH for the old halo clusters (70 objects); YH for the young halo clusters (30 objects); SG for clusters belonging to the Sagittarius dwarf (6 objects); and UN for clusters whose membership type remains unknown due to insufficient or unclassifiable data (7 objects).

In the following two sections, we consider some of the properties of these cluster sub-systems which were not considered by Mackey & Gilmore, but which nevertheless provide useful insights into the origin and evolution of objects in each of the different classes of Galactic globular cluster.

3 PROPERTIES OF THE DIFFERENT GALACTIC CLUSTER SUB-SYSTEMS

It is useful to briefly summarize the results of Mackey & Gilmore (2004), who examined some key properties of the Galactic globular

* HB-type is parametrized by the dimensionless morphology index $\frac{(B-R)}{(B+V+R)}$ (Lee 1990; Lee et al. 1994), where B is the number of HB stars which lie to the blue of the instability strip; R is the number of HB stars falling to the red of the instability strip; and V is the number of variable HB stars

† Mackey & Gilmore (2004) used the oldest isochrone from the theoretical HB models of Rey et al. (2001) as their fiducial

Table 1. Data and classifications for Galactic globular clusters.

Cluster Name	R_{gc} (kpc)	M_V	R_h (pc)	R_t (pc)	$\log I_h$ ($L_\odot \text{ pc}^{-2}$)	ϵ ($1 - b/a$)	[Fe/H]	HB-index	Class
NGC 104	7.4	-9.42	3.65	56.10	3.77	0.09	-0.76	-0.99	BD
NGC 288	12.0	-6.74	5.68	33.12	2.32	...	-1.24	0.98	OH
NGC 362	9.4	-8.41	2.00	39.83	3.89	0.01	-1.16	-0.87	YH
NGC 1261	18.2	-7.81	3.58	34.73	3.15	0.07	-1.35	-0.71	YH
Pal. 1	17.0	-2.47	2.16	28.41	1.45	0.22	-0.60	-1.00	BD
AM-1	123.2	-4.71	17.73	68.08	0.52	...	-1.80	-0.93	YH
Eridanus	95.2	-5.14	10.50	83.17	1.14	...	-1.46	-1.00	YH
Pal. 2	35.4	-8.01	5.38	54.27	2.87	0.05	-1.30	-0.10	YH
NGC 1851	16.7	-8.33	1.83	41.18	3.94	0.05	-1.22	-0.32	OH
NGC 1904	18.8	-7.86	3.00	31.30	3.32	0.01	-1.57	0.89	OH
NGC 2298	15.7	-6.30	2.43	20.17	2.88	0.08	-1.85	0.93	OH
NGC 2419	91.5	-9.58	17.88	214.07	2.46	0.03	-2.12	0.86	OH
Pyxis	41.7	-5.75	15.59	71.37	1.04	...	-1.30	-1.00	YH
NGC 2808	11.1	-9.39	2.12	43.42	4.23	0.12	-1.15	-0.49	OH
E3	7.6	-2.77	2.58	13.16	1.41	...	-0.80	...	UN
Pal. 3	95.9	-5.70	17.80	129.70	0.91	...	-1.66	-0.50	YH
NGC 3201	8.9	-7.46	3.90	41.38	2.93	0.12	-1.58	0.08	YH
Pal. 4	111.8	-6.02	17.15	105.78	1.07	...	-1.48	-1.00	YH
NGC 4147	21.3	-6.16	2.41	35.43	2.83	0.08	-1.83	0.66	SG
NGC 4372	7.1	-7.77	6.58	58.75	2.60	0.15	-2.09	1.00	OH
Rup. 106	18.5	-6.35	6.78	31.02	2.01	...	-1.67	-0.82	YH
NGC 4590	10.1	-7.35	4.60	90.02	2.74	0.05	-2.06	0.17	YH
NGC 4833	7.0	-8.16	4.56	33.75	3.08	0.07	-1.80	0.93	OH
NGC 5024	18.3	-8.70	5.75	112.62	3.09	0.01	-1.99	0.81	OH
NGC 5053	16.9	-6.72	16.70	65.21	1.37	0.21	-2.29	0.52	YH
NGC 5139	6.4	-10.29	6.44	87.92	3.63	0.17	-1.62	...	UN
NGC 5272	12.2	-8.93	3.39	115.53	3.64	0.04	-1.57	0.08	YH
NGC 5286	8.4	-8.61	2.21	26.75	3.89	0.12	-1.67	0.80	OH
AM-4	25.5	-1.60	3.65	11.48	0.65	...	-2.00	...	UN
NGC 5466	16.2	-6.96	10.41	158.36	1.88	0.11	-2.22	0.58	YH
NGC 5634	21.2	-7.69	3.96	61.28	3.01	0.02	-1.88	0.91	OH
NGC 5694	29.1	-7.81	3.33	43.30	3.21	0.04	-1.86	1.00	OH
IC 4499	15.7	-7.33	8.25	67.90	2.23	0.08	-1.60	0.11	YH
NGC 5824	25.8	-8.84	3.35	144.28	3.62	0.03	-1.85	0.79	OH
Pal. 5	18.6	-5.17	19.98	109.87	0.60	...	-1.41	-0.40	YH
NGC 5897	7.3	-7.21	7.61	43.54	2.25	0.08	-1.80	0.86	OH
NGC 5904	6.2	-8.81	4.60	61.96	3.33	0.14	-1.27	0.31	OH
NGC 5927	4.5	-7.80	2.54	36.88	3.44	0.04	-0.37	-1.00	BD
NGC 5946	5.8	-7.20	2.13	74.09	3.35	0.16	-1.38	0.69	OH
BH 176	9.7	-4.35	3.92	28.16	1.68	...	0.10	-1.00	BD
NGC 5986	4.8	-8.44	3.18	31.83	3.50	0.06	-1.58	0.97	OH
Lyngå 7	4.2	...	...	...	...	...	-0.62	-1.00	BD
Pal. 14	69.0	-4.73	24.72	113.07	0.24	...	-1.52	-1.00	YH
NGC 6093	3.8	-8.23	1.89	38.63	3.87	...	-1.75	0.93	OH
NGC 6121	5.9	-7.20	2.34	20.79	3.27	...	-1.20	-0.06	OH
NGC 6101	11.1	-6.91	7.61	32.36	2.13	0.05	-1.82	0.84	OH
NGC 6144	2.6	-6.75	4.01	82.21	2.62	0.25	-1.75	1.00	OH
NGC 6139	3.6	-8.36	2.41	25.03	3.71	0.05	-1.68	0.91	OH
Terzan 3	2.4	-4.61	2.84	12.89	2.07	...	-0.73	-1.00	BD
NGC 6171	3.3	-7.13	5.03	32.47	2.58	0.02	-1.04	-0.73	OH
1636-283	2.0	-3.97	1.30	7.84	2.49	...	-1.50	-0.40	YH
NGC 6205	8.7	-8.70	3.34	56.40	3.56	0.11	-1.54	0.97	OH
NGC 6229	29.7	-8.05	3.27	47.58	3.32	0.05	-1.43	0.24	YH
NGC 6218	4.5	-7.32	3.08	25.09	3.08	0.04	-1.48	0.97	OH
NGC 6235	4.1	-6.44	2.79	25.24	2.81	0.13	-1.40	0.89	OH
NGC 6254	4.6	-7.48	2.32	27.49	3.39	...	-1.52	0.98	OH
NGC 6256	1.8	-6.52	2.08	18.55	3.10	...	-0.70	-1.00	BD
Pal. 15	37.9	-5.49	15.70	64.74	0.93	...	-1.90	1.00	OH
NGC 6266	1.7	-9.19	2.47	18.00	4.02	0.01	-1.29	0.32	OH
NGC 6273	1.6	-9.18	3.13	36.27	3.81	0.27	-1.68	0.96	OH
NGC 6284	7.6	-7.97	3.47	102.72	3.24	0.03	-1.32	0.88	OH
NGC 6287	2.1	-7.36	2.03	28.43	3.46	0.13	-2.05	0.98	OH

Table 1. continued...

Cluster Name	R_{gc} (kpc)	M_V	R_h (pc)	R_t (pc)	$\log I_h$ ($L_\odot \text{ pc}^{-2}$)	ϵ ($1 - b/a$)	[Fe/H]	HB-index	Class
NGC 6293	1.4	-7.77	2.33	36.43	3.50	0.03	-1.92	0.90	OH
NGC 6304	2.2	-7.32	2.46	23.13	3.28	0.02	-0.59	-1.00	BD
NGC 6316	3.2	-8.35	2.27	18.97	3.76	0.04	-0.55	-1.00	BD
NGC 6341	9.6	-8.20	2.60	36.18	3.58	0.10	-2.28	0.91	OH
NGC 6325	1.1	-6.95	2.19	22.08	3.23	0.12	-1.17	0.84	OH
NGC 6333	1.7	-7.94	2.18	18.75	3.63	0.04	-1.75	0.87	OH
NGC 6342	1.7	-6.44	2.20	37.17	3.02	0.18	-0.65	-1.00	BD
NGC 6356	7.6	-8.52	3.27	35.24	3.51	0.03	-0.50	-1.00	BD
NGC 6355	1.8	-8.08	2.40	41.95	3.60	...	-1.50	0.62	OH
NGC 6352	3.3	-6.48	3.32	17.43	2.68	0.07	-0.70	-1.00	BD
IC 1257	17.9	-6.15	3.17	62.57	2.59	...	-1.70	1.00	OH
Terzan 2	0.9	-5.27	3.85	25.61	2.07	...	-0.40	-1.00	BD
NGC 6366	5.0	-5.77	2.75	15.92	2.56	0.16	-0.82	-0.97	OH
Terzan 4	1.3	-6.09	0.56	4.31	4.07	...	-1.60	1.00	OH
HP 1	6.1	-6.44	6.20	33.71	2.12	...	-1.55	0.75	OH
NGC 6362	5.1	-6.94	4.82	36.85	2.54	0.07	-0.95	-0.58	OH
Liller 1	1.8	-7.63	1.26	35.10	3.98	...	0.22	-1.00	BD
NGC 6380	3.2	-7.46	2.33	37.54	3.38	...	-0.50	-1.00	BD
Terzan 1	2.5	-4.90	6.22	18.03	1.50	...	-1.30	-1.00	YH
Ton. 2	1.4	-6.14	2.54	25.33	2.78	...	-0.50	-1.00	BD
NGC 6388	3.2	-9.42	1.95	18.06	4.32	0.01	-0.60	-1.00	BD
NGC 6402	4.1	-9.12	3.49	89.92	3.69	0.11	-1.39	0.65	OH
NGC 6401	2.7	-7.90	5.83	36.96	2.76	0.15	-0.98	0.35	OH
NGC 6397	6.0	-6.63	1.56	10.58	3.40	0.07	-1.95	0.98	OH
Pal. 6	2.2	-6.81	1.82	14.35	3.33	...	-1.09	-1.00	YH
NGC 6426	14.6	-6.69	5.78	79.66	2.28	0.15	-2.26	0.58	YH
Djorg. 1	4.1	-6.26	4.40	34.98	2.35	...	-2.00	...	UN
Terzan 5	2.4	-7.87	2.49	39.76	3.49	...	0.00	-1.00	BD
NGC 6440	1.3	-8.75	1.42	15.42	4.33	0.01	-0.34	-1.00	BD
NGC 6441	3.9	-9.64	2.18	27.23	4.31	0.02	-0.53	-1.00	BD
Terzan 6	1.6	-7.67	1.22	48.06	4.03	...	-0.50	-1.00	BD
NGC 6453	1.8	-6.88	1.03	60.04	3.86	0.09	-1.53	0.84	OH
UKS 1	0.8	-6.88	2.08	45.29	3.25	...	-0.50	-1.00	BD
NGC 6496	4.3	-7.23	6.26	17.63	2.43	0.16	-0.64	-1.00	BD
Terzan 9	1.6	-3.85	1.47	15.54	2.34	...	-2.00	0.25	YH
Djorg. 2	1.4	-6.98	1.62	20.52	3.50	...	-0.50	-1.00	BD
NGC 6517	4.3	-8.28	1.95	12.88	3.86	0.06	-1.37	0.62	OH
Terzan 10	2.4	-6.31	...	...	...	...	-0.70	-1.00	BD
NGC 6522	0.6	-7.67	2.36	37.30	3.45	0.06	-1.44	0.71	OH
NGC 6535	3.9	-4.75	1.52	16.54	2.67	0.08	-1.80	1.00	OH
NGC 6528	0.6	-6.56	0.99	38.08	3.76	0.11	-0.04	-1.00	BD
NGC 6539	3.1	-8.30	4.08	52.44	3.23	0.08	-0.66	-1.00	BD
NGC 6540	4.4	-5.38	0.26	10.21	4.45	...	-1.20	0.30	OH
NGC 6544	5.3	-6.66	1.39	1.61	3.51	0.22	-1.56	1.00	OH
NGC 6541	2.2	-8.37	2.42	60.27	3.71	0.12	-1.83	1.00	OH
2MASS-GC01	4.5	...	...	...	...	...	-1.20	...	UN
ESO280-SC06	14.3	...	...	...	...	...	-2.00	...	UN
NGC 6553	2.2	-7.77	2.71	14.24	3.37	0.17	-0.21	-1.00	BD
2MASS-GC02	4.1	...	...	...	...	...	...	...	UN
NGC 6558	1.0	-6.46	3.47	22.47	2.63	...	-1.44	0.70	OH
IC 1276	3.7	-6.67	3.69	32.96	2.66	...	-0.73	-1.00	BD
Terzan 12	3.4	-4.14	1.17	4.33	2.65	...	-0.50	-1.00	BD
NGC 6569	2.9	-8.30	4.14	21.63	3.22	...	-0.86	-0.82	OH
NGC 6584	7.0	-7.68	3.12	36.52	3.21	0.03	-1.49	-0.15	YH
NGC 6624	1.2	-7.49	1.88	47.22	3.58	0.06	-0.44	-1.00	BD
NGC 6626	2.7	-8.18	2.54	18.36	3.59	0.16	-1.45	0.90	OH
NGC 6638	2.3	-7.13	1.84	18.51	3.45	0.01	-0.99	-0.30	OH
NGC 6637	1.9	-7.64	2.20	22.10	3.50	0.01	-0.70	-1.00	BD
NGC 6642	1.7	-6.77	1.78	24.61	3.34	0.03	-1.35	-0.04	YH
NGC 6652	2.8	-6.68	1.91	13.16	3.24	0.20	-0.96	-1.00	OH
NGC 6656	4.9	-8.50	3.03	26.97	3.57	0.14	-1.64	0.91	OH
Pal. 8	5.6	-5.52	2.14	50.85	2.68	...	-0.48	-1.00	BD

Table 1. continued...

Cluster Name	R_{gc} (kpc)	M_V	R_h (pc)	R_t (pc)	$\log I_h$ ($L_\odot \text{ pc}^{-2}$)	ϵ ($1 - b/a$)	[Fe/H]	HB-index	Class
NGC 6681	2.1	-7.11	2.43	20.71	3.20	0.01	-1.51	0.96	OH
NGC 6712	3.5	-7.50	2.75	14.93	3.25	0.11	-1.01	-0.62	OH
NGC 6715	19.2	-10.01	3.82	58.23	3.97	0.06	-1.79	0.54	SG
NGC 6717	2.4	-5.66	1.40	20.38	3.10	0.01	-1.29	0.98	OH
NGC 6723	2.6	-7.84	4.07	26.60	3.05	...	-1.12	-0.08	OH
NGC 6749	5.0	-6.70	2.53	11.97	3.00	...	-1.60	1.00	OH
NGC 6752	5.2	-7.73	2.72	64.39	3.35	0.04	-1.56	1.00	OH
NGC 6760	4.8	-7.86	4.69	27.90	2.93	0.04	-0.52	-1.00	BD
NGC 6779	9.7	-7.38	3.41	25.15	3.02	0.03	-1.94	0.98	OH
Terzan 7	16.0	-5.05	6.55	49.06	1.52	...	-0.82	-1.00	SG
Pal. 10	6.4	-5.79	1.70	5.29	2.98	...	-0.10	-1.00	BD
Arp 2	21.4	-5.29	15.89	105.24	0.84	...	-1.84	0.53	SG
NGC 6809	3.9	-7.55	4.46	25.10	2.85	0.02	-1.81	0.87	OH
Terzan 8	19.1	-5.05	7.56	30.25	1.39	...	-1.99	1.00	SG
Pal. 11	7.9	-6.86	5.63	37.13	2.37	...	-0.39	-1.00	BD
NGC 6838	6.7	-5.60	1.92	10.43	2.80	...	-0.73	-1.00	BD
NGC 6864	14.6	-8.55	2.83	43.84	3.65	0.07	-1.16	-0.07	OH
NGC 6934	12.8	-7.46	2.74	38.23	3.24	0.01	-1.54	0.25	YH
NGC 6981	12.9	-7.04	4.35	45.25	2.67	0.02	-1.40	0.14	YH
NGC 7006	38.8	-7.68	4.59	76.54	2.88	0.01	-1.63	-0.28	YH
NGC 7078	10.4	-9.17	3.18	64.42	3.79	0.05	-2.26	0.67	YH
NGC 7089	10.4	-9.02	3.11	71.75	3.75	0.11	-1.62	0.92	OH
NGC 7099	7.1	-7.43	2.68	42.68	3.25	0.01	-2.12	0.89	OH
Pal. 12	15.9	-4.48	7.11	96.78	1.22	...	-0.94	-1.00	SG
Pal. 13	26.7	-3.74	3.45	23.34	1.55	...	-1.74	-0.20	YH
NGC 7492	24.9	-5.77	9.16	62.67	1.51	0.24	-1.51	0.81	OH

cluster sub-systems. They found that the majority of globular clusters associated with nearby dwarf galaxies are, in terms of HB morphology and core-radius distribution, essentially indistinguishable from those in the Galactic YH class. Furthermore, the YH objects are observed to be characterized by large, energetic orbits around the Galaxy. These orbits are generally of high eccentricity and intermediate inclination, and cover a very large range in orbital angular momentum, including highly retrograde orbits. A small fraction of OH clusters have similar orbits, but most have much smaller energies and eccentricities. Many BD clusters (unsurprisingly) have disk-like motions, and/or orbits which are confined to the Galactic bulge (e.g., Dinescu et al. (2003)).

Mackey & Gilmore also examined the distribution of cluster ages in each of the sub-systems, using data from the relative age study of Salaris & Weiss (2002). These authors determined relative ages for 55 globular clusters using several different CMD-based techniques. A significant fraction of the YH clusters are ~ 3 Gyr younger than the majority of the OH clusters (see Fig. 9 in Mackey & Gilmore (2004)). However, several OH clusters are also seen to be somewhat younger, while a number of YH clusters appear to be just as old as the oldest measured OH clusters. The latter result is interesting because it hints explicitly at the possible existence of a third parameter, in addition to [Fe/H] and age, which governs HB morphology in a cluster.

Finally, Mackey & Gilmore observed that the YH clusters follow a considerably steeper age-metallicity relationship than do most of the OH and BD clusters (Mackey & Gilmore, Fig. 10). This shows that YH-type clusters apparently underwent much slower evolution to higher metallicities than did the OH and BD objects.

Taken together, the results of Mackey & Gilmore (2004) fully

support the conclusion of Zinn (1993) that the YH clusters are not native members of the Galaxy, but rather are of external origin. Furthermore, Mackey & Gilmore (2004) showed that some 15 per cent of the OH clusters possess core radii, orbital motions, ages and HB morphologies which suggest that they too are of external origin. The remainder of the OH group, together with the BD system, appear to have been formed in the (proto) Galaxy. In the following sub-sections, we consider some additional properties of the three cluster sub-systems, focusing on structural parameters such as R_h and ellipticity, as well as on cluster luminosities and surface brightnesses.

3.1 Distribution with Galactocentric radius

It has long been known that YH-type clusters typically reside at much larger Galactocentric distances than do most OH and BD clusters (see e.g., Zinn (1993); van den Bergh (1993); Mackey & Gilmore (2004)). This is clearly seen in Fig. 1, which shows the cumulative distribution in the number of globulars as a function of R_{gc} . Each of the three globular cluster sub-systems are quite distinct in terms of their distribution of Galactocentric radii. We can quantify the significance of this statement using simple Kolmogorov-Smirnov (K-S) tests on each pair of distributions. These show that there is only a 2 per cent chance that the BD and OH samples were drawn from the same parent population. The probability that the YH clusters share the same parent distribution as either the OH or BD clusters is even smaller: 0.015 per cent, and 0.01 per cent, respectively.

The metal-rich BD clusters are, as expected, the most centrally concentrated system, exhibiting a sharp cut-off at $R_{gc} = 7.9$ kpc.




Figure 1. Cumulative distributions in Galactocentric radii R_{gc} for the three Galactic globular cluster sub-systems.

Only two BD clusters lie outside this radius – BH 176 ($R_{gc} = 9.7$ kpc) and Pal. 1 ($R_{gc} = 17$ kpc). It is interesting to note that both of these clusters have been tentatively associated with the candidate dwarf galaxy in Canis Major, and/or the Monoceros tidal stream (see e.g., Frinchaboy et al. (2004)). In addition, both van den Bergh & Mackey (2004; hereafter Paper I) and Mackey & Gilmore (2004) have previously noted that Pal. 1 has an unusually high metallicity for its Galactocentric radius. In this respect it strongly resembles the Sagittarius clusters Ter. 7 and Pal. 12 (see e.g., Fig. 8 in Mackey & Gilmore (2004)). Furthermore, examining Table 1, Pal. 1 and BH 176 are, respectively, the lowest luminosity ($M_V = -2.5$) and the third lowest luminosity ($M_V = -4.4$) clusters in the BD ensemble. This adds additional weight to the suspicion that these two objects might not be native BD clusters.

The OH clusters are considerably more extended in R_{gc} than the BD objects, and dominate the inner Galactic halo. However, they are not nearly as extended as the YH clusters, which dominate the outer halo. Only six of the 70 OH clusters (9 per cent) lie beyond $R_{gc} = 20$ kpc, and of these, only one (the very unusual cluster NGC 2419 – see Section 4.3) has $R_{gc} > 40$ kpc. In contrast, one third of the YH system (10 clusters) have $R_{gc} > 20$ kpc, and six of these objects fall outside $R_{gc} = 40$ kpc. The most distant YH cluster is AM-1 at $R_{gc} \sim 120$ kpc. As noted by Mackey & Gilmore (2004), the new classifications presented in Table 1 show that the YH clusters also extend into even central regions of the Galaxy. These objects are therefore not exclusively a remote-halo population.

3.2 Distribution of cluster luminosities

In Fig. 2 we plot histograms of the luminosity distributions for clusters in each of the three sub-systems. From this plot it is seen that both the BD and OH systems have peaks near $M_V \sim -7.75$. Furthermore, these two systems contain very few low luminosity clusters. Only 11 per cent of BD clusters and 4 per cent of OH




Figure 2. Luminosity distributions for the Galactic globular cluster sub-systems.

clusters are fainter than $M_V = -5.5$. This contrasts with the situation for the YH system, which contains a significant population of low luminosity objects. It is seen that 27 per cent of the clusters in this sub-system are fainter than $M_V = -5.5$. In addition, the distribution of YH clusters peaks at somewhat lower luminosity ($M_V \sim -7$) than do the OH and BD distributions. Again, we can evaluate the significance of the differences between the three distributions using K-S tests. These show that there is only a 5 per cent chance that the OH and YH clusters share the same parent luminosity distribution. However, the K-S tests are unable to discriminate between the OH and BD distributions, and the YH and BD distributions – the results do not confirm statistically significant differences between these parent populations, but they also do not confirm statistically significant similarities. This result is at least partly due to the relatively small BD and YH sample sizes. As discussed above, the primary difference between the distributions is the presence (or not) of a small percentage of low luminosity clusters. Hence, it would be necessary for the sample sizes to be larger to confirm (or deny) this difference at a significant level. This is, of course, not possible, since we included all known Galactic globular clusters in Table 1.

We can, however, clearly state that the luminosity distribution of YH clusters is significantly broader than that for the OH clusters. Combined with our observations from Section 3.1, this result is similar to that obtained by Kavelaars & Hanes (1997), who demonstrated that outer halo globular clusters (i.e., those metal-poor clusters with $R_{gc} > 8$ kpc) have a broader luminosity function than the inner halo globular clusters (i.e., those metal-poor clusters with $R_{gc} < 8$ kpc). The present result is, however, distinct from this, as we have not used R_{gc} as a discriminator – indeed, Fig. 1 shows that there is a significant population of OH clusters with $R_{gc} > 8$ kpc, and vice versa for YH clusters. Our observations imply that LF broadness is intrinsic to cluster sub-population rather than simply location within the Galaxy.

As discussed by Kavelaars & Hanes (1997), it seems likely

that this observed difference in broadness, as well as the apparent dearth of low luminosity clusters in the OH and BD systems, can (at least in part) be attributed to the destructive influence of tidal forces and bulge and disk shocks in the inner Galaxy. A number of authors (see e.g., Fall & Zhang (2001)) have shown that gravitational shocks, in addition to cluster relaxation and evaporation, are important in shaping the globular cluster mass function. Since the YH clusters typically lie at larger Galactocentric radii than do OH and BD clusters, they should be much less strongly affected by destructive external forces.

In addition, if the accretion hypothesis is correct, many of the YH clusters have spent some fraction of their life outside the Galaxy in a more benign tidal environment. Consider, for example, the clusters associated with the Sagittarius dwarf. While it is at present very difficult to constrain the early orbital evolution of this galaxy, models by Jiang & Binney (2000) have shown that its Galactocentric distance at apocentre ~ 11 Gyr ago most likely fell in the range $60 < R_{gc} < 250$ kpc, with the exact value depending on its initial mass (specifically, the amount of dark matter it contained). Models by Helmi & White (2001) and Ibata et al. (2001), which assume comparatively small initial masses, show that Sagittarius may have followed a similar orbit to that presently observed (which has apocentric radius ~ 65 kpc, pericentric radius ~ 15 kpc, and an orbital period ~ 1 Gyr) for the last $\sim 10 - 12$ Gyr. In each of these scenarios, the Sagittarius dwarf, and its remaining globular clusters, spend a significant fraction of their lives away from the inner regions of the Galaxy. The observation that four of the Sagittarius clusters are still associated with the main body of this galaxy provides further evidence that these clusters have not suffered prolonged tidal stresses due to the Galaxy. Tidal forces strong enough to disrupt a globular cluster are likely to be so strong that they would also pull that cluster out of its parent dwarf galaxy.

Together the factors described above may help to explain why the YH system has a greater fraction of low luminosity clusters than do the OH and BD systems. If destructive effects are indeed responsible, then the YH luminosity distribution may well indicate that globular clusters are formed with a broad LF. Alternatively, as suggested by Kavelaars & Hanes (1997), the shape of the cluster LF may be dependent on the environment in which clusters form.

Finally, it is interesting to examine the combined luminosity distribution of globular clusters in the four nearby dwarf galaxies studied by Mackey & Gilmore (2004). This distribution is plotted in Fig. 3, and bears a strong resemblance to the luminosity distribution of the YH clusters. A significant fraction of the external globular clusters (25 per cent) are fainter than $M_V = -5.5$, and in fact this fraction is almost identical to that observed for the YH clusters. A K-S test gives a 92 per cent chance that the YH and external clusters have the same parent luminosity distribution, but only a 3 per cent chance that the OH and external clusters share the same parent luminosity distribution. This observation adds another example to the list of characteristics in which the YH sub-system closely resembles the group of globular clusters associated with nearby Local Group dwarf galaxies (Mackey & Gilmore 2004). It therefore offers further evidence in favour of the hypothesis that much, or all, of the ensemble of YH clusters is of external origin.

3.3 Distribution of half-light and tidal radii

Figs. 4 and 5 show the distributions of half-light radii (R_h) and tidal radii (R_t), respectively, for the three Galactic cluster sub-systems. Just as Mackey & Gilmore (2004) demonstrated for core radii, it is clear from these two Figures that the majority of diffuse clusters –




Figure 3. Combined luminosity distribution for the globular clusters in the LMC, SMC and Fornax and Sagittarius dwarf galaxies. Note how similar this distribution is to that for the YH clusters (Fig. 2).

that is, clusters with large half-light and tidal radii – fall into the YH category. Conversely, the large majority of OH and BD objects are seen to be compact. Once again, we apply K-S tests to determine the significance of the differences between the three distributions. These calculations show that there is a less than 1 per cent chance that the OH and YH distributions in R_h were drawn from the same parent population, and a less than 0.2 per cent chance that the BD and YH distributions in R_h share the same parent population. For the BD and OH R_h distributions, this probability is 8 per cent. Exactly similar results are obtained for the R_t distributions.

Given what is known about the distributions in Galactocentric radius for the three sub-systems (Section 3.1), this result is not unexpected when considering R_t . The tidal radii of clusters are determined by how deeply their orbits dive into the Galactic potential in which they reside (see e.g., King (1962)). In fact, it has been shown that R_t is most strongly correlated with the distance of a cluster at the pericentre of its orbit (van den Bergh 1994). Hence, since the majority of clusters which reside at large R_{gc} belong to the YH ensemble, it is to be expected that this group also contains the largest fraction of clusters with large R_t . By the same token, the BD clusters are the most centrally concentrated in R_{gc} , and consequently all the clusters in this sub-sample have comparatively small tidal radii.

The distributions in R_h are somewhat more intriguing. It has long been known that the half-light radii of Galactic globular clusters correlate with their Galactocentric distances, in the sense that the most remote clusters are also the largest (e.g., van den Bergh & Morbey (1984)). Hence, with only this fact in mind, one might expect the YH system to possess most of the clusters with large R_h . This is indeed what is observed. However, the fact that the YH clusters are strongly suspected of being accreted in a number of minor merger events presents a challenge for this explanation. This is so because, for clusters which have been more-or-less randomly accreted, there is no reason to expect half-light radii to follow the same trend with R_{gc} as that which is observed for clusters which are native to the Galaxy. We address this puzzle in more detail in Section 4.2.

Irrespective of the above result, the distributions in R_h are interesting because, as described in Section 1, a cluster's half-light radius is stable over many relaxation times. Hence, the observed distribution of half-light radii for the YH clusters should, to a large extent, reflect the distribution of radii with which these clusters




Figure 4. Distributions of half-light radii (R_h) for the Galactic globular cluster sub-systems.




Figure 5. Distributions of tidal radii (R_t) for the Galactic globular cluster sub-systems.

were formed. If this is the case, then the fact that the OH and BD systems contain almost no diffuse clusters with $R_h > 7$ pc allows two possibilities: either (i) the distributions in R_h for these two ensembles also remain mostly unmodified from their original states, and the dearth of diffuse objects is due to environmental influence on cluster formation (i.e., no diffuse clusters were formed); or (ii) the native Galactic cluster population has been strongly modified by destructive tidal forces. From the cluster destruction studies

of Gnedin & Ostriker (1997) and Dinescu, Girard, & van Altena (1999) it is clear that destructive influences, such as evaporation due to two-body relaxation as well as disk and bulge shocks, must be at least partly responsible. These most strongly affect diffuse clusters – that is, objects with large R_h and low M_V . For example, in the formalism of Dinescu et al. (1999) the destruction rate due to both bulge and disk shocks is directly proportional to R_h^3 , and inversely proportional to cluster mass. Furthermore, clusters with short orbital periods and small pericentric radii (i.e., many OH clusters and most BD clusters) are most strongly affected. In their study, Dinescu et al. estimated destruction rates for two of our BD clusters, 25 of our OH clusters, and 10 of our YH clusters. They provide a list of the nine clusters with the largest destruction rates (these objects have expected survival times of ~ 10 Gyr or less). Of these, one is a BD cluster, 7 are OH clusters, and one a YH cluster. Clearly the native population is most strongly affected. In addition, several of the nine have large R_h (e.g., NGC 288, Pal. 5, and NGC 6144), while two (NGC 6121 and 6397) have been shown to have depleted luminosity functions at low stellar masses (Kanas et al. 1995; Piotto et al. 1997). This is expected in a cluster if it has been subjected to strong external tidal forces.

Assuming that some fraction of the native Galactic clusters has been destroyed as discussed above, it is possible to place constraints on the initial population. Examining Fig. 4, one sees that 33 per cent of YH clusters have $R_h > 7$ pc. Mackey & Gilmore (2004) concluded that there are presently ~ 100 native Galactic members in the BD/OH set (recall that ~ 15 per cent of the OH clusters are possibly of external origin). If the initial distribution of cluster sizes for native Galactic globulars resembled that presently observed for YH objects, then this population constitutes only 67 per cent of the original population, since essentially no OH or BD clusters are observed to have $R_h > 7$ pc. Hence the original population of native Galactic globular clusters may have numbered ~ 150 objects. This result is consistent with both the estimate of Mackey & Gilmore (2004) who postulated an original Galactic population numbering ~ 200 objects, and the result of Gnedin & Ostriker (1997) who concluded that up to half of the present population of globular clusters are likely to be destroyed during the next Hubble time.

It is important to point out that the above argument explicitly depends on the assumption that the OH and BD systems originally had R_h distributions similar to that which is presently observed for YH clusters. Alternatively we again note the possibility that formation conditions did not allow the production of diffuse clusters in the inner Galaxy. In that case our estimate of ~ 150 original Galactic globular clusters is an over-estimate. However, not enough information about ancient cluster formation conditions is available at present to allow this possibility to be confirmed or rejected.

3.4 Distribution of surface brightnesses

In Fig. 6 we plot the distributions of I_h , the half-light intensity (essentially equivalent to the mean surface brightness within R_h), for the three Galactic globular cluster sub-systems. Since this quantity is closely related to both R_h and M_V , it is not surprising to see that the vast majority of low surface brightness clusters belong to the YH sub-system. The distribution of I_h for this ensemble is quite broad with no prominent maximum. In this respect it differs significantly from the much sharper singly-peaked distributions that are observed for the BD and OH clusters. A Kolmogorov-Smirnov test shows that there is only a 1 per cent chance that the YH and BD samples were drawn from the same parent distribution, and a




Figure 6. Distributions of half-light intensity (I_h) for the three Galactic cluster sub-systems.

less than 0.1 per cent probability that the YH and OH samples were drawn from the same parent distribution.

The lack of low surface brightness clusters among the OH and BD systems results in a scarcity of clusters with $I_h < 100 L_\odot \text{ pc}^{-2}$ and $R_{gc} < 15 \text{ kpc}$ (as can be seen by inspecting Table 1). It is also of interest to note that all the nine known Galactic globular clusters with $I_h < 20 L_\odot \text{ pc}^{-2}$ are metal-poor objects having $[\text{Fe}/\text{H}] < -1.2$. As was already pointed out in Sections 3.2 and 3.3, these effects are likely due to the tidal destruction of fragile low surface brightness clusters that ventured too close to the massive bulge of the Galaxy. An additional complication is that observational selection effects militate against the discovery of low surface brightness clusters against the rich foreground of stars in the direction of the Galactic bulge.

It is interesting to note that the location of the peak surface brightness for the OH and BD clusters coincides closely with the location of the (marginal) surface brightness peak for the YH clusters, at $\log I_h \sim 3.25$. As previously noted, the YH population has apparently not been as strongly affected by tidal forces as have the OH and BD populations. This sub-system may therefore provide interesting constraints on the initial distribution of cluster surface brightnesses (although, as noted previously, I_h is more sensitive to cluster evolution than is R_h , because the evaporation of stars, along with stellar evolution, causes variations in M_V). Thirty-seven per cent of YH clusters have $I_h < 100 L_\odot \text{ pc}^{-2}$, while for the BD and OH clusters this fraction is essentially zero. If we again set the number of native Galactic members in the BD and OH groups to be ~ 100 then, assuming the distribution of surface brightnesses for this ensemble used to resemble that now observed for the YH clusters, the present population of native Galactic clusters represents 63 per cent of the original population, which must therefore have numbered ~ 160 clusters. This is consistent with the estimate already given in the previous Section. It is interesting to note that in a thought experiment where we assume all YH clusters with $I_h < 100 \text{ mag pc}^{-2}$ to have been destroyed, the amplitude of the




Figure 7. Distributions of ellipticity for the three Galactic globular cluster sub-systems.

surface brightness peak at $\log I_h \sim 3.25$ rises to ~ 0.32 – which is similar to the amplitudes of the peaks that are observed for the OH and BD clusters.

3.5 Cluster ellipticities

Next we consider the ellipticity distributions of Galactic globular clusters, which are plotted in Fig. 7. This Figure shows some interesting features. In particular, the YH clusters appear on average to be somewhat less flattened than the OH clusters, although the small YH sample precludes this difference being confirmed as significant by a K-S test. Additional measurements will be required to further investigate this possible difference. If the average flattening of YH clusters does turn out to be less than that of OH clusters, this would be unexpected for several reasons. Frenk & Fall (1982) found the oldest LMC globular clusters to be slightly *more* elliptical than typical Galactic globular clusters. A similar result was obtained by Goodwin (1997). Since a number of other parameters show a strong similarity between the YH clusters and the sample of external clusters (of which LMC objects make up a significant fraction) it is perhaps surprising that the YH clusters should be less elliptical on average than the OH clusters. Frenk & Fall (1982) also found that older LMC clusters are typically less elliptical than younger LMC clusters. If this were true in general, one would have expected the YH clusters, many of which are $\sim 3 \text{ Gyr}$ younger than most of the OH clusters, to be more elliptical than these Galactic natives.

Goodwin (1997) has argued that the strength of the tidal field in which a cluster evolves is the predominant factor in shaping its ellipticity. This is so because a strong tidal field is quite effective in reducing the triaxiality of a cluster and in removing any angular momentum it may have, thus reducing its ellipticity. This may explain why clusters in the LMC are more elliptical, on average, than Galactic clusters. However, this prediction appears to be at odds with the present observation that the YH clusters, which generally

reside at large R_{gc} and have supposedly spent at least some fraction of their lifetimes in low-mass galaxies, may typically be less flattened than are the OH and BD clusters, which reside in a strong tidal field. One possible explanation for this puzzle is that recent bulge and disk shocks have re-introduced velocity anisotropy into many of the OH and BD clusters (Goodwin 1997). If so, this might explain the tail to high ellipticities for the OH clusters, which is not present for the YH clusters.

Finally, it is interesting to note that all three of the lowest surface brightness ($I_h < 30 L_\odot \text{pc}^{-2}$) clusters for which ellipticity values have been published are unusually flattened objects with $1 - b/a > 0.2$. These clusters are NGC 5053, NGC 7492, and Pal. 1. Examination of Digitized Sky Survey (DSS) images confirms this unusually high ellipticity for NGC 5053 and 7492 (the DSS image of Pal. 1 is too faint for useful measurements), so this result does not seem to be a measurement artifact due to low surface brightness. Clearly, more observational data would be valuable. Such information could be obtained in the near future because there are still a significant number of low surface brightness clusters for which ellipticity measurements are not yet available. In this connection, it is of interest to note that the very faint ($M_V = -3.0_{-0.7}^{+2.0}$) and large ($R_h = 23 \pm 10$ pc) globular cluster-like object recently discovered in the outer halo of the Galaxy by Willman et al. (2004) is of exceedingly low half-light intensity ($I_h = 0.40_{-0.37}^{+2.00} L_\odot \text{pc}^{-2}$ – i.e., $\log I_h = -0.4_{-1.1}^{+0.8}$), and also appears to be significantly flattened in its outer regions.

If the observed ellipticity of the lowest surface brightness globulars were due to rotation, then this would hint at a physical relation between rotation and surface brightness. Alternatively, it is possible that some globular clusters might be embedded in small dark matter halos so that the shapes of the lowest surface brightness clusters reflect the axis ratios of their dark matter mini-halos. In this connection it is noted that Mashchenko & Sills (2004a; 2004b) have recently investigated the formation and evolution of globular clusters in different types of dark matter mini-halos. They conclude that the presence of obvious tidal tails is the only observational evidence that can reliably rule out significant amounts of dark matter in a globular cluster. Furthermore, they write that “*cosmological dark matter halos are known to have noticeably non-spherical shapes; a stellar cluster relaxing inside such a halo would have a spherical distribution in its denser part where stars dominate dark matter, and would exhibit isodensity contours of increasingly larger eccentricity in its outskirts where DM becomes the dominant mass component.*” It is currently believed (e.g., Bailin & Steinmetz (2004)) that massive cosmological dark halos are triaxial with c/a ratios of ~ 0.6 . However, the numerical resolution on cosmological N -body simulations is presently insufficient to reliably predict the likely flattenings of subhalos with masses that are only a millionth of those of the massive halos themselves. Furthermore, the shapes of such dark matter subhalos orbiting the Galaxy may have been strongly affected by tidal stripping, which rapidly reduces the masses of dark matter subhalos once they have been accreted onto larger structures (e.g., Gao et al. (2004)). Even so, the case for a detailed new systematic study of globular cluster ellipticities is strong, especially because of the present availability of sensitive wide field cameras at a number of large telescopes. Such a study should certainly include low surface brightness clusters, because such objects are still very poorly studied. It would be particularly interesting to examine how ellipticity changes with projected radius, i.e. do clusters get more or less flattened with increasing projected radius?




Figure 8. Plot of $\log R_h$ vs. M_V for Galactic globular clusters with $R_{gc} > 15$ kpc (filled circles), and those in the LMC (open squares), SMC (star), and Fornax and Sagittarius dwarf galaxies (open triangles, and crosses, respectively). The best fit line, Eq. 1 is marked by the solid line, while the two 1σ error lines are marked by the dotted lines.

4 THE LOG R_H VERSUS M_V PLANE

4.1 The link between cluster size and luminosity

It is informative to consider the correlation between cluster luminosity and half-light radius. Several authors have examined this relationship in the past, arriving at different conclusions. For example, Ostriker & Gnedin (1997) found R_h to be strongly correlated with luminosity[†] for clusters with $5 < R_{gc} < 60$ kpc, such that $R_h \propto (L/L_\odot)^{-0.63}$ with correlation coefficient 0.77. However, van den Bergh, Morbey & Pazder (1991) found that half-light radius also correlates quite strongly with Galactocentric radius R_{gc} for clusters with $R_{gc} < 40$ kpc: $R_h \propto R_{gc}^{0.5}$ with correlation coefficient $0.37_{-0.19}^{+0.16}$. They found that if this dependence is taken into account, cluster size appears uncorrelated with luminosity – that is, a “normalized” radius R_h^* , which is the size a cluster would have at given R_{gc} , does not correlate with luminosity. A similar conclusion was reached by McLaughlin (2000) who found that for the full sample of Galactic globular clusters, a weak dependence of R_h on luminosity disappears when the variation of R_h on R_{gc} (McLaughlin found $R_h \propto R_{gc}^{0.4}$) is taken into account. However, McLaughlin stresses that he cannot reproduce the result of Ostriker & Gnedin (1997) for clusters with $5 < R_{gc} < 60$ kpc.

It seems possible that these different conclusions are due, at least in part, to different sample selections. McLaughlin (2000) considered the full Galactic globular cluster system, and the sample of van den Bergh et al. (1991) contained 98 clusters with $R_{gc} < 40$ kpc. In contrast, the ensemble of Ostriker & Gnedin (1997) consisted of 58 non-core-collapsed clusters with $5 < R_{gc} < 60$ kpc. In Section 3, we discussed the idea that the globular cluster sample

[†] assuming a roughly constant M/L ratio for Galactic globular clusters

consisting of objects in the central Galaxy has had its properties strongly modified by external (tidal) influences. We found support for this notion, in agreement with a number of previous authors. With this in mind, in the present context it makes sense to consider a cluster sample which is as free from tidal modification as possible. To this end, we consider a Galactic globular cluster sample consisting only of objects with $R_{gc} > 15$ kpc. The majority of clusters in this ensemble are YH members. In Paper I we made plots of $\log R_h$ versus M_V for both this Galactic sample, as well as for a sample consisting of globular clusters in the LMC, SMC, and Fornax and Sagittarius dwarf galaxies (Figs. 6 & 7 of Paper I). Interestingly, both of these plots appear to show a strong inverse correlation between cluster size and luminosity, particularly for clusters in the external dwarf galaxies. We reproduce these data here, in Fig. 8.

Apart from four notable outliers, the majority of clusters in Fig. 8 exhibit a clear trend in R_h with M_V . Three of the deviant clusters – Pal. 1, AM-4, and Pal. 13 – are compact but of very low luminosity, and lie below and to the right of the majority of the data. All three of these objects have been shown to be extremely poorly populated globular clusters – AM-4 does not even possess any RGB or HB stars (Inman & Carney 1987). Pal. 1 has been studied by Rosenberg et al. (1998), who measured an unusually flat mass function, very different from those measured for 21 more luminous Galactic globular clusters. These authors concluded that Pal. 1 has most likely been modified by tidal forces in combination with the evaporation of cluster members. Assuming this to be the case, they find Pal. 1 must presently be very close to its final destruction. Similarly, Siegel et al. (2001) showed that Pal. 13 has been strongly affected by tidal forces and is on the threshold of dissolution, a result supported by the unexpectedly high internal velocity dispersion measured for this object by Côté et al. (2002). We note that the inner halo cluster E3 also inhabits this region of the $\log R_h$ vs. M_V plane. This cluster is also exceedingly sparse. The photometric study by McClure et al. (1985) found a sharp drop-off in the cluster luminosity function several magnitudes below the main sequence turn off. This supports the suggestion by van den Bergh et al. (1980) that E3 has been strongly modified by evaporative processes.

Since Pal. 1, AM-4 and Pal. 13 have apparently been strongly affected by external destructive forces, we exclude them from our present sample and analysis. All three clusters are apparently very close to dissolution, so the lower right hand corner of Fig. 8 may constitute the ultimate “graveyard” of compact globular clusters. This leaves the anomalous object NGC 2419, which is the fourth deviant cluster in Fig. 8, and which is discussed in more detail in Section 4.3, below. We note that the inclusion or omission of this object from the sample does not strongly affect our results (below), although the formal errors are reduced if NGC 2419 is not included.

Applying a least-squares analysis to the sample, one obtains a best-fit line:

$$\log R_h = 0.25M_V + 2.5, \quad (1)$$

where the formal errors are ± 0.03 on the slope and ± 0.18 on the intercept. This line is plotted in Fig. 8, along with the 1σ error lines. These errors represent the case when NGC 2419 is not included in the analysis. The trend defined by Eq. 1 is strong, with correlation coefficient 0.83. It is important to note however, that the distribution of cluster radii in Fig. 8 is dominated by scatter.

If one replaces the integrated magnitude in Eq. 1 with $M_V = -2.5 \log(L/L_\odot) + M_{V,\odot}$ one finds that $R_h \propto (L/L_\odot)^{-0.625}$, which is almost exactly the result of Ostriker & Gnedin (1997) if

M/L is assumed to be approximately constant for Galactic globular clusters. The physical significance of Eq. 1 is not clear, however. One possibility is that the relationship between R_h and M_V has been imposed purely through the dynamical evolution of clusters. If this is the case, then the majority of evolution must have been in M_V as R_h is stable over cluster lifetimes (see Section 1). M_V fades as a cluster evolves, due to both stellar evolution and the evaporation of cluster members. The rate of fading due to stellar evolution is regulated by the cluster IMF (because more massive stars die earlier), while evaporation is caused by both external influences (e.g., tidal stripping) and internal relaxation. Loosely bound clusters are susceptible to both processes (see e.g., Gnedin & Ostriker (1997)), although we note that our sample has been chosen to minimise possible modifications due to tidal shocking. In contrast, compact clusters are significantly more stable, although very compact objects suffer accelerated evaporation due to internal relaxation (see e.g., Figs. 6 & 7 in Gnedin & Ostriker (1997)). Given the complexity of these combined factors, detailed N -body modelling will be required in order to assess how significant evolutionary effects are on the R_h vs. M_V diagram, how they vary with cluster concentration, and whether they can explain the observed correlation.

An alternative possibility is that this correlation represents some signature of globular cluster formation. If this is so, the details of this signature are not immediately evident, as dynamical evolution (e.g., fading in M_V) must again be taken into account if the presently observed trend is to be traced back to its original form. In considering whether Eq. 1 is at least partly due to cluster formation conditions, it is important to note that a number of previous authors (e.g., van den Bergh et al. (1991); McLaughlin (2000)) have claimed that a dependence of the half-light radius R_h on Galactocentric distance R_{gc} might contribute to the observed correlation between cluster size and luminosity for Galactic globular clusters. A physical interpretation of this argument is that the properties of the Galactic globular cluster system were set by the cluster formation process, which is environment-dependent – that is, the parameters with which a cluster was formed were modulated by forces that depend on Galactocentric radius (see e.g., McLaughlin (2000) for discussion). The basis for this idea can be seen in Figure 9, which shows a plot of R_h as a function of R_{gc} for the full sample of Galactic globular clusters. There is a clear trend of increasing R_h with increasing R_{gc} . A formal fit to these data yields

$$\log R_h = 0.42 \log R_{gc} + 0.19, \quad (2)$$

where the errors are ± 0.04 on both the slope and intercept, and the correlation coefficient is 0.65. This result is consistent with those found previously by both McLaughlin (2000) and van den Bergh et al. (1991).

In our present sample however, the extent to which a correlation between R_h and Galactocentric radius can drive that between R_h and M_V is not clear. All our Galactic clusters are in the outer halo, with $R_{gc} > 15$ kpc. Fig. 9 shows that these objects have a very large scatter in R_h , and furthermore that diffuse clusters with $R_{gc} \sim 15$ kpc do *not* have systematically smaller R_h than diffuse clusters with $R_{gc} \sim 100$ kpc. Any apparent trend in R_h with R_{gc} for these outer halo clusters is most likely caused by the surprising *lack* of compact clusters at very large Galactocentric radii (see Section 4.2 below). In addition, the majority of the sample with $R_{gc} > 15$ kpc are YH members, and as such are very possibly accreted objects. In this scenario, such clusters would not be expected to exhibit any trend between R_h and R_{gc} amongst themselves. We also note that if the data in Fig. 8 are split into the Galactic and “external” clusters, then fitting these two groups separately repro-




Figure 9. Plot of $\log R_h$ vs. $\log R_{gc}$ for all Galactic globular clusters. The BD clusters are marked by open triangles; the OH clusters by crosses; the YH clusters by filled circles; the Sagittarius clusters by open circles; and the unclassified (UN) clusters by open squares. The vertical dotted line indicates $R_{gc} = 15$ kpc.

duces Eq. 1 for each to within 1σ , albeit both with lower significance (due to the smaller number of points in each group). It is curious that the globular clusters in the LMC, SMC, and Fornax and Sagittarius dSph galaxies follow such a similar trend in Fig. 8 to the outer halo Galactic clusters, even though they are members of significantly different galaxies and have evolved in different environments. Even so, it is observed that in the LMC, the most diffuse clusters (Reticulum, ESO121-SC03, NGC 1841, 2257) all lie at large galactocentric radii, as does the very extended cluster 1 in the Fornax dSph, so there seems to be at least a weak dependence of R_h on radius in these galaxies.

Ultimately, the most important conclusion to draw from Figures 8 and 9 is that the Galactic globular clusters with large R_h are both faint, and occur at large Galactocentric radii. With present data, it is not possible to identify for certain the origin of the observed trend in R_h with M_V . Although there is some evidence that a correlation between R_h and R_{gc} could be at least partly responsible, at present an intrinsic relationship between R_h and M_V , perhaps a relic of the cluster formation process, cannot and should not be ruled out. This puzzle seems well suited for study using large-scale realistic N -body simulations (see e.g., Aarseth (1999)), which with the advent of powerful dedicated hardware (e.g., GRAPE-6 – see Makino et al. (2003)) are now nearly capable of direct globular cluster modelling.

4.2 Where are the compact outer halo globular clusters?

As noted above, examining Fig. 9 in more detail reveals a surprising result: no compact YH clusters occur at $R_{gc} > 40$ kpc. This observation is intriguing because the presently accepted paradigm is that the young halo objects have been accreted by the Galaxy via minor mergers. Their R_h values should therefore be indepen-




Figure 10. Plot of $\log I_h$ vs. $\log R_{gc}$ for all Galactic globular clusters. The point styles are as in Fig. 9.

dent of R_{gc} . This is so because the chaotic acquisition of clusters is expected to result in a wide spread of cluster sizes across a broad range of R_{gc} values. Indeed, this is exactly what is seen at intermediate Galactocentric radii ($10 < R_{gc} < 40$ kpc). The dispersion in R_h for the YH clusters at such radii is considerably larger than that for the clusters that are intrinsically associated with the Milky Way, and in fact closely matches the spread in R_h among the Sagittarius clusters. At very small R_{gc} we only really see compact YH clusters. However, recalling the apparent influence tidal forces have had on the OH and BD samples, this is not unexpected because extended clusters are especially vulnerable to rapid tidal destruction in the inner Galaxy.

It is therefore the handful of YH clusters at very large Galactocentric radii[§] ($R_{gc} > 40$ kpc) which are the most intriguing. A simple examination of Fig. 4 shows that, among a randomly selected sample of six YH clusters, one should expect approximately three to have $R_h < 5$ pc, or more than four to have $R_h < 10$ pc. However, Fig. 9 shows that there are *no* compact clusters among the six observed at very large Galactocentric radii. If the very outer young halo clusters originally belonged to now-defunct dwarf galaxies, as the presently accepted paradigm would seem to suggest, then there is no clear reason why their dispersion in R_h should not be similar to that observed for YH clusters with $10 < R_{gc} < 40$ kpc.

Inspection of Table 1 reveals an additional interesting fact. All of these six outer YH clusters are of relatively low luminosity (indeed, this result might be expected given Fig. 8 and Eq. 1). The surface brightness I_h is derived from the parameters R_h and M_V , and is therefore useful in the present analysis. Fig. 10 shows a plot of I_h as a function of R_{gc} for all Galactic globular clusters. In the inner Galaxy, there is no significant trend in I_h with R_{gc} , but (as discussed in Section 3.4) there is a marked lack of low surface brightness clusters – which are the most difficult to find against the back-

[§] AM-1, Eridanus, Pyxis, Pal. 3, Pal. 4, and Pal. 14

drop of the Galactic bulge, and which are most vulnerable to tidal disruption. At intermediate R_{gc} , the YH clusters are seen to have a significantly larger spread in I_h than do the native Galactic clusters. In fact, the dispersion in I_h observed at these Galactocentric radii for the YH clusters matches closely the spread in I_h exhibited by the Sagittarius objects. At $R_{gc} > 40$ kpc only very low surface brightness clusters are found, i.e. only low luminosity clusters with large R_h . Apart from NGC 2419, which is also interesting for other reasons (see Section 4.3, below), *there are no globular clusters of even average surface brightness in the outer Galactic halo.*

We can think of several possible scenarios that might account for this lack of compact/luminous clusters at very large R_{gc} :

(1) Clusters with $R_{gc} > 40$ kpc once belonged to one, or more likely several, now-destroyed parent dwarf galaxies which disintegrated at large R_{gc} and which *only* possessed diffuse clusters at the time of merger.

(2) Clusters with $R_{gc} > 40$ kpc were outlying objects in one, or more likely several, parent dwarf galaxies, and were lost very early on during merger events which subsequently proceeded at smaller R_{gc} .

(3) Clusters with $R_{gc} > 40$ kpc never belonged to a parent dwarf galaxy and were formed as isolated objects at large Galactocentric radii.

Since little is known about the properties of galaxies which might have merged with ours at early epochs, it is not possible to definitively rule out any of these options. *If* these galaxies were similar to the Fornax and Sagittarius dSph galaxies observed today, then options 1 and 2 seem implausible. One could perhaps imagine a scenario in which any compact cluster members of a dwarf galaxy were formed close to the centre of the parent galaxy and soon merged into its core, leaving only diffuse clusters. However, both the Fornax and Sagittarius dwarf galaxies contain clusters with a wide range of R_h , so, unless the hypothetical parent dwarf galaxies of the outer halo globulars were radically different to Fornax and Sagittarius, option 1 appears unlikely. In addition, Fornax is at a distance of $R_{gc} \sim 140$ kpc and shows no signs of being strongly tidally influenced by the Galaxy, let alone disrupted. It also possesses two outlying clusters (Fornax 1 and 5) which have not been removed. Furthermore, the Sagittarius dwarf is *much* closer to the Galactic centre ($R_{gc} \sim 20$ kpc), and still apparently possesses four clusters, of which two are diffuse objects (Terzan 8 and Arp 2). Hence, option 2 also appears unlikely in this scenario.

Having outlined this argument, we point out that we cannot be certain that the galaxies which merged with ours at early epochs were in any way similar to Fornax and Sagittarius. Indeed, studies of the chemical abundance patterns in halo stars and in local dSph galaxies suggest that perhaps they were not (see Mackey & Gilmore (2004) for a brief summary). It is possible that the hypothetical parent dwarf galaxies of the outer YH clusters, along with their member clusters, were systematically more diffuse than Fornax and Sagittarius and their clusters. If this were the case, it might explain why these diffuse galaxies are now destroyed whereas Fornax and Sagittarius are still (mostly) intact. In this scenario, we cannot rule out options 1 and 2.

Option 3 also seems plausible, as the outermost YH clusters might help form the upper end of a trend in increasing R_h with increasing R_{gc} if considered together with the BD and OH clusters (Fig. 9). However, at least four of these outer young halo clusters are believed to be several Gyr younger than the oldest Galactic globulars (e.g., Stetson et al. (1999); Sarajedini (1997)). This raises the question: why (and how) did cluster formation occur at such re-




Figure 11. Plot of $\log R_h$ vs. M_V for all Galactic globular clusters. The point styles are again as in Fig. 9. The giant M31 globular cluster G1 is marked by two linked stars – these represent the two discrepant measurements of R_h for this cluster. The upper point is the measurement of Meylan et al. (2001), while the lower point is that of Barmby et al. (2002).

note locations *after* it occurred deeper within the Galactic potential well. Furthermore, it is not clear why (or how) such cluster formation would produce only diffuse, low luminosity clusters, and why these objects are similar in so many ways to those found in local dwarf galaxies.

The outermost clusters are clearly very important tracers of how remote parts of the Galactic halo were formed or assembled, as well as the properties of the systems in which they might have formed. It is therefore clearly of great interest to study them in considerably more detail than has been achieved to date.

4.3 The structures of very luminous clusters

Figure 11 shows a plot of $\log R_h$ vs. M_V for all Galactic globular clusters. The most important thing to note from this Figure is that the upper left envelope of the main distribution of globular clusters has a rather sharp edge. Only a very few clusters lie a significant distance above this upper edge. If we shift the line defined by Eq. 1 upwards so that it has intercept 2.95 (as plotted on Fig. 11) then one finds that the line

$$\log R_h = 0.25 M_V + 2.95 \quad (3)$$

closely describes the upper envelope of the cluster distribution. Only three Galactic globular clusters lie above this line: ω Centauri, M54, and NGC 2419. A number of authors, such as Bekki & Freeman (2003), Tsuchiya et al. (2003), and Ideta & Makino (2004) (and references therein), have demonstrated that the observed properties of ω Centauri are consistent with it being the stripped core of a former dwarf spheroidal or dwarf elliptical galaxy. Furthermore, Ibata, Gilmore & Irwin (1994; 1995) have shown that M54 (= NGC 6715) is associated with the central region of the Sagittarius dwarf galaxy, and some authors (see e.g. Layden & Sarajedini (2000))

have argued that M54 may actually be the core of this galaxy. Finally, in Paper I we suggested that NGC 2419 might also be the remaining core of a defunct dwarf spheroidal-type galaxy. In this connection it would clearly be of great interest to check if NGC 2419, like ω Centauri (e.g., Bedin et al. (2004); Hilker et al. (2004); and references therein), also shows evidence for age/metallicity substructure in its color-magnitude diagram. However, evidence for such substructure in NGC 2419 might be difficult to obtain because the mean metallicity ($\langle[\text{Fe}/\text{H}]\rangle = -2.12$) of NGC 2419 is so much lower than that of ω Centauri ($\langle[\text{Fe}/\text{H}]\rangle = -1.62$). As a result the expected range in metallicities for the population components of NGC 2419 is expected to be quite small. NGC 2419 is also significantly more distant than ω Centauri ($R_{\odot} = 84.2$ kpc), further complicating any observations.

It is intriguing that ω Centauri, M54, and NGC 2419, which have all independently been previously suggested as the cores of now-disrupted dwarf galaxies, should constitute a group of objects which have such strikingly different structures than other luminous Galactic clusters. This observation suggests a similar origin for these three objects, since each has clearly evolved under quite different circumstances and in different environments. Inspection of Table 1 shows that three out of the four Galactic globular clusters brighter than $M_V = -9.5$ fall above Eq. 3. The lone exception is NGC 6441 ($M_V = -9.64$, $R_h = 2.2$ pc). The high metallicity ($[\text{Fe}/\text{H}] = -0.53$) of this cluster, which is located at $R_{gc} = 3.9$ kpc, militates in favor of its assignment to the population associated with the Galactic bulge or inner disk. It is noted in passing that the apparently normal globular NGC 5053 also falls very close to (but still marginally below) the line defined by Eq. 3.

Since the majority of the most luminous Galactic globular clusters are now suspected of being the cores of former dwarf galaxies it seems likely that the same might also be true in the globular cluster system associated with M31. This suspicion is strengthened by the observation that the cluster M31 Mayall No. II (= G1), which may be the most luminous globular cluster associated with the Andromeda galaxy, has $M_V = -10.94$ and $R_h = 14$ pc (Meylan et al. 2001), although Barmby, Holland, & Huchra (2002) find $R_h = 3.1$ pc (using the distance of Meylan et al. (2001)). Adopting either of these R_h measurements, G1 also falls above the line defined by Eq. 3, i.e. in the same region as ω Cen, M54, and NGC 2419. In this connection, we note that Meylan et al. (2001) found their HST/WFPC2 photometry of G1 to be consistent with an internal metallicity spread of ~ 0.5 dex. Furthermore Bekki & Chiba (2004) have recently argued that the observed properties of G1 can be explained if this object is the tidally stripped nucleus of a former dwarf elliptical (dE,N) galaxy.

Finally, Harris et al. (2002) constructed a catalogue of structural measurements for globular clusters in NGC 5128 (= Centaurus A). The 14 most luminous of these have recently been studied spectroscopically by Martini & Ho (2004), who hypothesize that some of these clusters are actually the stripped nuclei of dwarf galaxies, based on their large masses and the possible detection of extra-tidal light by Harris et al. (2002) for several. On the assumption that our line Eq. 3 provides a reasonable means of discriminating objects which may be the cores of former dwarf galaxies, the speculation of Martini & Ho (2004) is supported by the observation that all 14 of the clusters in their sample lie above and to the left of this line. In other words, all 14 have structures and luminosities more similar to those of ω Cen, M54, NGC 2419, and G1 than to those of the majority of “ordinary” globular clusters associated with our Galaxy. Harris, Harris & Geisler (2004) have recently found that NGC 5128 contains 980 ± 120 globulars, so that potential dwarf

galaxy remnants represent at least ~ 1.4 per cent of the clusters associated with this galaxy.

5 CONCLUSIONS

In terms of their observed characteristics (metallicities, HB morphologies, structures, shapes, luminosities, and surface brightnesses) the Galactic globular clusters, considered as a whole, constitute a relatively inhomogeneous class of objects. From a physical point of view, it therefore makes sense to try to identify subgroups made up of clusters with similar characteristics. One of the most successful classification schemes for the Galactic globular clusters is that introduced by Zinn (1993) and van den Bergh (1993) (see also the references therein), whereby clusters are segregated according to their metallicity and HB morphology. Recently this scheme has been updated to cover the entire Galactic globular cluster system by Mackey & Gilmore (2004).

In the present paper, we have presented the complete list of cluster classifications of Mackey & Gilmore (2004). Several objects in the Galactic halo possess properties which are consistent with the hypothesis that they are the stripped cores of former dwarf galaxies (see below). The remaining majority of “normal” globular clusters fall into three major categories – bulge/disk (BD), “old” halo (OH), and “young” halo (YH) clusters. As recently re-emphasized by Mackey & Gilmore (2004) (see also Section 3, above), the majority of BD and OH objects seem to have formed in the (proto) Galaxy, while the YH objects possess characteristics which indicate that they most likely formed in extra-Galactic systems and were accreted into the Galactic halo during subsequent merger events.

We have examined several of the properties of these three sub-systems which were not considered by Mackey & Gilmore (2004). The BD, OH, and YH clusters typically occupy different regions of the Galaxy. The metal-rich BD objects are the most centrally concentrated, exhibiting a sharp cut-off in their Galactocentric radial distribution at $R_{gc} = 7.9$ kpc. In contrast, the OH clusters dominate the inner halo, with the majority residing within $R_{gc} = 20$ kpc, while the YH clusters typically occupy the outer halo, with more than half lying at $R_{gc} > 15$ kpc.

In terms of cluster structures, luminosities, surface brightnesses, and ellipticities, the YH clusters are clearly distinct from the OH and BD objects. The OH and BD populations have apparently been strongly modified by destructive tidal forces and bulge and disk shocks, so that these sub-systems possess few extended clusters, few low luminosity clusters, and few low surface brightness clusters. In comparison, the YH group contains a significant fraction of extended clusters with half-light radii $R_h > 7$ pc, and low luminosity clusters with $M_V > -5.5$. The luminosity distribution of YH clusters matches closely that observed for clusters in four nearby dwarf galaxies (the LMC, SMC, and Fornax and Sagittarius dwarf spheroidals), and as such, offers further evidence for the external origin of much or all of the YH ensemble. By examining how the observed distributions of half-light radii and mean surface brightnesses for the YH clusters would have to be modified to match those observed for native Galactic globulars, we estimate that the original population of Galactic globular clusters may have numbered $\sim 150 - 160$ objects. This result is consistent with several previous estimates in the literature.

The distribution of cluster ellipticities shows that the YH clusters are, on average, less flattened than are the OH and BD clusters. A possible explanation of this result is that recent bulge and disk shocks have introduced significant velocity anisotropy (and

hence ellipticity) into many BD and OH clusters (Goodwin 1997). We also note that all three of the lowest surface brightness clusters for which ellipticity values have been published are unusually flattened objects, as is the newly discovered very low surface brightness object of Willman et al. (2004). This may indicate an intrinsic relationship between cluster rotation and surface brightness. An alternative, and perhaps more attractive, explanation invokes the influence of dark matter. In the globulars of lowest surface density the structure of the potential energy well might be dominated by a triaxial dark matter mini-halo (Mashchenko & Sills 2004a; Mashchenko & Sills 2004b), rather than by the baryonic material associated with Population II stars. The flattening of the isophotes of low surface brightness globular clusters might therefore provide information on their (still hypothetical) dark matter mini-halos. We briefly made the case for a comprehensive study of Galactic globular cluster ellipticities.

We have also investigated the relationship between cluster luminosity and half-light radius. A strong correlation between the two is observed for clusters in the outer Galactic halo and also among clusters in the LMC, SMC, and Fornax and Sagittarius dwarf galaxies. Several authors have previously asserted that any correlation between R_h and M_V for Galactic globular clusters can be explained by the fact that R_h also appears to correlate with Galactocentric radius. However, this cannot explain why the outer halo Galactic clusters, which are mostly YH objects, follow the trend in R_h with M_V . If these objects are accreted, no trend in R_h with R_{gc} is expected.

An additional puzzling observation is the lack of compact globular clusters in the remote halo. We observe six YH clusters with $R_{gc} > 40$ kpc; however all are diffuse and of low luminosity. Based on the observed distributions of R_h and M_V for YH clusters, one might have expected several of the six to be compact and of greater than average luminosity. We discussed several scenarios which might explain the dearth of compact clusters with very large R_{gc} ; however none appear particularly attractive.

Finally, if all the Galactic globular clusters are plotted on the $\log R_h$ vs. M_V plane, several clusters stand well clear of the main body of objects. At very low luminosities we observe a number of compact objects (Pal. 1, Pal. 13, E3 and AM-4) which have apparently been strongly affected by shrinkage due to cooling and evaporation of cluster stars, and stripping by Galactic tides. The most interesting result, however, concerns the upper envelope of the main distribution of clusters, which is sharply defined. Only three Galactic globulars lie significantly above a line (Eq. 3) describing this upper envelope – ω Cen, M54, and NGC 2419. All three of these objects have previously, and independently, been hypothesized as being the stripped cores of former dwarf spheroidal-type galaxies. This suggests that the position of a cluster in the $\log R_h$ vs. M_V plane provides a powerful tool for the segregation between “normal” globular clusters and those which might previously have been associated with the cores of dwarf galaxies. This suspicion is reinforced by the additional observation that the massive cluster G1 in M31, which has been previously suggested to be the remaining core of a nucleated dwarf elliptical galaxy, also lies above Eq. 3, as do 14 of the most luminous clusters in NGC 5128, which have again been postulated as constituting the remnant cores of now defunct dwarf galaxies.

ACKNOWLEDGEMENTS

We thank Peter Stetson for commenting on an early draft of this paper, and George Wallerstein for correspondence about the color-magnitude diagrams of dwarf spheroidals. We are also indebted to Marcin Sawicki, Liang Gao and Julio Navarro for helpful discussions, and we acknowledge the anonymous referee, whose thoughtful and constructive comments greatly improved this work. ADM recognises financial support from PPARC in the form of a Postdoctoral Fellowship.

REFERENCES

- Aarseth S. J., 1999, *Celest. Mech. Dynam. Astron.*, 73, 127
- Aarseth S. J., Heggie D. C., 1998, *MNRAS*, 297, 794
- Bailin J., Steinmetz M., 2004, *ApJ*, submitted (astro-ph/0408163)
- Barmby P., Holland S., Huchra J. P., 2002, *AJ*, 123, 1937
- Bedin L. R., Piotto G., Anderson J., King I. R., Cassisi S., Momany Y., 2004, *Mem. Soc. Astr. It.*, in press (astro-ph/0406076)
- Bekki K., Freeman K. C., 2003, *MNRAS*, 346, L11
- Bekki K., Chiba M., 2004, *A&A*, 417, 437
- Côté P., 1999, *AJ*, 118, 406
- Côté P., Djorgovski S. G., Meylan G., Castro S., McCarthy J. K., 2002, *ApJ*, 574, 783
- Davidge T. J., 2000, *ApJS*, 126, 105
- Dinescu D. I., Girard T. M., van Altena W. F., 1999, *AJ*, 117, 1792
- Dinescu D. I., Girard T. M., van Altena W. F., López C. E., 2003, *AJ*, 125, 1373
- Fall S. M., Zhang Q., 2001, *ApJ*, 561, 751
- Frenk C. S., Fall S. M., 1982, *MNRAS*, 199, 565
- Frinchaboy P. M., Majewski S. R., Crane J. D., Reid I. N., Rocha-Pinto H. J., Phelps R. L., Patterson R. J., Muñoz R. R., 2004, *ApJ*, 602, L21
- Gao L., De Lucia G., White S. D. M., Jenkins A., 2004, *MNRAS*, 352, L1
- Goodwin S. P., 1997, *MNRAS*, 286, 39
- Gnedin O. Y., Ostriker J. P., 1997, *ApJ*, 474, 223
- Harris G. L. H., Harris W. E., Geisler D., 2004, *AJ*, 128, 723
- Harris W. E., 1996, *AJ*, 112, 1487
- Harris W. E., Harris G. L. H., Holland S. T., McLaughlin D. E., 2002, *AJ*, 124, 1435
- Helmi A., White S. D. M., 2001, *MNRAS*, 323, 529
- Hilker M., Kayser A., Richtler T., Willemsen P., 2004, *A&A*, 422, 9
- Ibata R. A., Gilmore G. F., Irwin M. J., 1994, *Nature*, 370, 194
- Ibata R. A., Gilmore G. F., Irwin M. J., 1995, *MNRAS*, 277, 781
- Ibata R. A., Lewis G. F., Irwin M., Totten E., Quinn T., 2001, *ApJ*, 551, 294
- Ideta M., Makino J., 2004, *ApJ*, in press (astro-ph/0408431)
- Inman R. T., Carney B. W., 1987, *AJ*, 93, 1166
- Jiang I. -G., Binney J., 2000, *MNRAS*, 314, 468
- Jordán A., 2004, *ApJ*, 613, L117
- Kanas I. N., Griffiths W. K., Dickens R. J., Penny A. J., 1995, *MNRAS*, 272, 265
- Kavelaars J. J., Hanes D. A., 1997, *MNRAS*, 285, L31
- King I., 1962, *AJ*, 67, 471
- Layden A. C., Sarajedini A., 2000, *AJ*, 119, 1760
- Lee Y. -W., 1990, *ApJ*, 363, 159
- Lee Y. -W., Demarque P., Zinn R., 1994, *ApJ*, 423, 248
- Lightman A. P., Shapiro S. L., 1978, *Rev Mod. Phys.* 50, 437
- Mackey A. D., Gilmore G. F., 2003a, *MNRAS*, 340, 175
- Mackey A. D., Gilmore G. F., 2003b, *MNRAS*, 343, 747
- Mackey A. D., Gilmore G. F., 2004, *MNRAS*, in press
- Makino J., Fukushige T., Koga M., Namura K., 2003, *PASJ*, 55, 1163
- Martini P., Ho L. C., 2004, *ApJ*, in press (astro-ph/0404003)
- Mashchenko S., Sills A., 2004a, *ApJ*, in press (astro-ph/0409605)
- Mashchenko S., Sills A., 2004b, *ApJ*, in press (astro-ph/0409606)
- McClure R. D., Hesser J. E., Stetson P. B., Stryker L. L., 1985, *PASP*, 97, 665
- McLaughlin D. E., 2000, *ApJ*, 539, 618

- Meylan G., Sarajedini A., Jablonka P., Djorgovski S. G., Bridges T., Rich R. M., 2001, *AJ*, 122, 830
- Ortolani S., Bica E., Barbuy B., 1997, *MNRAS*, 284, 692
- Ostriker J. P., Gnedin O. Y., 1997, *ApJ*, 487, 667
- Piotto G., Cool A. M., King I. R., 1997, *AJ*, 113, 1345
- Rey S. -C., Yoon S. -J., Lee Y. -W., Chaboyer B., Sarajedini A., 2001, *AJ*, 122, 3219
- Rosenberg A., Saviane I., Piotto G., Aparicio A., Zaggia S. R., 1998, *AJ*, 115, 648
- Salaris M., Weiss A., 2002, *A&A*, 388, 492
- Sarajedini A., 1997, *AJ*, 113, 682
- Siegel M. H., Majewski S. R., Cudworth K. M., Takamiya M., 2001, *AJ*, 121, 935
- Spitzer L., Thuan T. X., 1972, *ApJ*, 175, 31
- Stetson P. B., et al., 1999, *AJ*, 117, 247
- Tsuchiya T., Dinescu D. I., Korchagin V. I., 2003, *ApJ*, 598, L29
- van den Bergh S., 1993, *ApJ*, 411, 178
- van den Bergh S., 1994, *AJ*, 108, 2145
- van den Bergh S., Morbey C. L., 1984, *Astronomy Express*, 1, 1
- van den Bergh S., Mackey A. D., 2004, *MNRAS*, 354, 713 (= Paper I)
- van den Bergh S., Demers S., Kunkel W. E., 1980, *ApJ*, 239, 112
- van den Bergh S., Morbey C., Pazder J., 1991, *ApJ*, 375, 594
- Webbink R. F., 1985, in Goodman J., Hut P., eds., *Proc. IAU Symp.* 113, *Dynamics of Star Clusters*. Kluwer, Dordrecht, p. 541
- Willman B., et al., 2004, *AJ*, submitted (astro-ph/0410416)
- Zinn R., 1993a, in Smith G. H., Brodie J. P., eds., *ASP Conf. Ser.* 48, *The Globular Cluster-Galaxy Connection*. Astron. Soc. Pac., San Francisco, p. 38

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.